

**CONSUMER INFORMATION**

**Pr** **PRO-FLUCONAZOLE**  
Fluconazole Tablets, House Standard

This leaflet is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about PRO-FLUCONAZOLE. Contact your doctor or pharmacist if you have any questions about the drug.

**ABOUT THIS MEDICATION****What the medication is used for:**

PRO-FLUCONAZOLE is one of a group of medicines called antifungals.

PRO-FLUCONAZOLE is prescribed by your doctor and is used to treat infections caused by fungi including yeasts and may also be used to stop you from getting a fungal infection. The most common cause of fungal infection is a yeast called *Candida*.

**What it does:**

Fluconazole selectively interferes with the normal sterol production in fungi and helps to stop fungal growth.

**When it should not be used:**

Do not take PRO-FLUCONAZOLE if you have ever had an allergic reaction to:

- any of the ingredients of PRO-FLUCONAZOLE (see What the non-medicinal ingredients are)
- other medicines you have taken to treat a fungal infection.

The symptoms may include itching, reddening of the skin or difficulty in breathing.

Do not take PRO-FLUCONAZOLE if you are taking any of the following drugs:

- Cisapride\* (used as a gastrointestinal motility agent),
- Terfenadine\* or astemizole\* (antihistamines for allergies),
- Erythromycin (an antibiotic for treating infections),
- Pimozide (for treating schizophrenia) and
- Quinidine (used for irregular heartbeats).

\* not marketed in Canada

**What the medicinal ingredient is:**

Fluconazole

**What the non-medicinal ingredients are:**

Croscarmellose sodium, dibasic calcium phosphate anhydrous, FD&C Red No. 3 and No. 40 Lake dye, magnesium stearate, microcrystalline cellulose and povidone.

**What dosage forms it comes in:**

**Tablets:** 100 mg

**WARNINGS AND PRECAUTIONS**

Tell your doctor if you become pregnant while taking PRO-FLUCONAZOLE.

**BEFORE** you use PRO-FLUCONAZOLE talk to your doctor or pharmacist if you:

- have liver or kidney problems
- are allergic to any other medicines including those used to treat yeast and other fungal infections
- have abnormal levels of potassium, calcium or magnesium in your blood
- are pregnant or are planning to become pregnant.
- are breast-feeding or planning to breast-feed. Fluconazole is excreted in human breast milk. Breast-feeding is not recommended.
- have heart disease such as heart conditions, blood disorders or any other medical conditions.
- are driving and using machines. It should be taken into account that occasionally dizziness or seizures may occur.
- are taking or have taken any other medicines, including medicines obtained without a prescription

**Use in Children**

Your doctor will decide whether this medication is suitable for your child.

**INTERACTIONS WITH THIS MEDICATION**

**Drugs that may interact with PRO-FLUCONAZOLE include:**

- Alfentanil, fentanyl or methadone (used to treat pain)
- Amitriptyline, nortriptyline (used to treat migraine and other conditions)
- Amphotericin B and Voriconazole (used to treat fungal infections)
- Benzodiazepines such as midazolam, triazolam, or similar medicines (used to help you sleep or for anxiety)
- Calcium channel blockers or losartan (for lowering blood pressure)
- Carbamazepine or phenytoin (used to control epilepsy)
- Celecoxib (used to treat some types of arthritis and certain other types of pain)
- Cimetidine (for heartburn and peptic ulcers)
- Coumarin-Type Anticoagulants (used to thin the blood to prevent blood clots)
- Cyclophosphamide, or vinca alkaloids (for treating some forms of cancer)
- Cyclosporine, sirolimus or tacrolimus (to prevent transplant rejection)
- Halofantrine (to treat malaria)
- HMG-CoA reductase inhibitors (statins) (for lowering cholesterol)
- Medicines for treating infections (antibiotics) such as azithromycin, erythromycin, rifampin or rifabutin

- Non-steroidal anti-inflammatory drugs (such as acetylsalicylic acid and ibuprofen) that are used to treat pain and fever
- Oral Contraceptives
- Prednisone (used to treat many types of inflammatory and allergic conditions)
- Saquinavir or zidovudine, also known as AZT (used in HIV-infected patients)
- Sulfonylureas and other Oral Hypoglycemics (medicines for diabetes)
- Theophylline (used to control asthma)
- Tofacitinib (used to treat rheumatoid arthritis)
- Vitamin A (as a trans-retinoid acid used to treat acne)
- Water tablets (diuretics), such as hydrochlorothiazide, (used to treat fluid retention and high blood pressure)

**PROPER USE OF THIS MEDICATION**

**Usual dose:**

Take PRO-FLUCONAZOLE only as directed by your doctor.

**Overdose:**

In case of drug overdose, contact a health care practitioner, hospital emergency department or regional Poison Control Centre immediately, even if there are no symptoms.

**Missed Dose:**

If a dose of this medication has been missed, it should be taken as soon as possible. However, if it is almost time for the next dose, skip the missed dose and go back to the regular dosing schedule. Do not double dose.

**SIDE EFFECTS AND WHAT TO DO ABOUT THEM**

Like all medicines, PRO-FLUCONAZOLE may cause some side effects.

The most common side effects are:

- Headache,
- Skin rash,
- Abdominal pain,
- Diarrhea,
- Nausea and vomiting.

**SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM**

| Symptom / effect   | Talk with your doctor or pharmacist |              | Stop taking drug and seek immediate emergency medical attention |
|--|-------------------------------------|--------------|---|
|  | Only if severe                      | In all cases |   |
| <b>Exfoliative skin disorders:</b><br>Severe skin reactions, such as a rash that causes blistering, itching all over the body, reddening of the skin or itchy red spots, swelling of eyelids, face or lips, peeling or lost skin |                                     | √            |   |
| <b>Hepatic necrosis</b><br>(death of liver cells which may cause abdominal pain and dark urine, fever, light-coloured stool, and jaundice (a yellow appearance to the skin and white portion of the eyes))                       |                                     | √            |   |
| <b>Heart conditions:</b><br>Unstable or irregular heartbeat (e.g., QT prolongation, <i>torsade de pointes</i> )  |                                     | √            |   |
| <b>Allergic Reaction</b><br>with symptom such as swelling of the face, throat, mouth, extremities, difficulty in breathing, rash or itching  |                                     |              | √   |

*This is not a complete list of side effects. For any unexpected effects while taking PRO-FLUCONAZOLE, contact your doctor or pharmacist immediately.*

**HOW TO STORE IT**

Store between 15°C and 30°C.

Keep out of reach and sight of children.

### **Reporting Side Effects**

You can help improve the safe use of health products for Canadians by reporting serious and unexpected side effects to Health Canada. Your report may help to identify new side effects and change the product safety information.

#### **3 ways to report:**

- Online at [MedEffect](#);
- By calling 1-866-234-2345 (toll-free);
- By completing a Consumer Side Effect Reporting Form and sending it by:
  - Fax to 1-866-678-6789 (toll-free), or
  - Mail to: Canada Vigilance Program  
Health Canada, Postal Locator 0701E  
Ottawa, ON  
K1A 0K9Postage paid labels and the Consumer Side Effect Reporting Form are available at [MedEffect](#).

*NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.*

### **MORE INFORMATION**

This document plus the full product monograph, prepared for health professionals, can be obtained by contacting Pro Doc Ltée at 1-800-361-8559, [www.prodoc.qc.ca](http://www.prodoc.qc.ca) or [info@prodoc.qc.ca](mailto:info@prodoc.qc.ca).

This leaflet was prepared by  
Pro Doc Ltée, Laval, Québec, H7L 3W9

Last revised: August 26, 2016