

**READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE
PATIENT MEDICATION INFORMATION**

**PRO-MIRTAZAPINE
Mirtazapine Tablets, USP**

Read this carefully before you start taking **PRO-MIRTAZAPINE** and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about **PRO-MIRTAZAPINE**.

What is PRO-MIRTAZAPINE used for?

- PRO-MIRTAZAPINE is used in adults to relieve symptoms of depression.

How does PRO-MIRTAZAPINE work?

PRO-MIRTAZAPINE belongs to a group of medicines called anti-depressants. The way PRO-MIRTAZAPINE works to treat depression is unknown. PRO-MIRTAZAPINE is thought to have an effect in the brain on chemicals called serotonin and norepinephrine.

What are the ingredients in PRO-MIRTAZAPINE?

Medicinal ingredients: Mirtazapine

Non-medicinal ingredients: Colloidal Silicon Dioxide, Iron Oxide Yellow, Lactose, Magnesium Stearate, Microcrystalline Cellulose, Polyethylene Glycol, Sodium Starch Glycolate, Titanium Dioxide, and the following:

15 mg tablets also contain: Hydroxypropyl Methylcellulose, Polysorbate 80

30 mg tablets also contain: Iron Oxide Red, Polyvinyl Alcohol, Talc

PRO-MIRTAZAPINE comes in the following dosage forms:

Tablets: 15 mg and 30 mg

Do not use PRO-MIRTAZAPINE if you are:

- allergic to mirtazapine or any of the other ingredients in the medication.
- currently taking or have recently taken monoamine oxidase (MAO) inhibitors (including some types of anti-depressant and anti-Parkinson treatments) in the past 14 days.

PRO-MIRTAZAPINE is not for use in patients under 18 years of age.

To help avoid side effects and ensure proper use, talk to your healthcare professional before you take PRO-MIRTAZAPINE. Talk about any health conditions or problems you may have, including if you:

- have ever shown unusual sensitivity (rash or any other signs of allergy) to any medication;
- have heart problems;
- have had a heart attack or stroke;
- have liver or kidney problems;
- are taking a medicine to lower your blood pressure;
- have a history of drug abuse;
- have a history of mania;
- have a history of seizures or fits;
- have diabetes;
- have prostate problems;
- have glaucoma or increased pressure in your eyes;

- have a history of suicidal behaviour or other mental health problems, such as schizophrenia and bipolar disorder (alternating periods of elation/overactivity and depressed mood);
 - have a history of high cholesterol and/or high triglycerides (fats in the blood);
 - are 65 years of age or older;
 - are lactose intolerant or have one of the following rare hereditary diseases:
 - Galactose intolerance
 - Lapp lactase deficiency
 - Glucose-galactose malabsorption
- Because lactose is a non-medicinal ingredient in PRO-MIRTAZAPINE.

Other warnings you should know about:

PRO-MIRTAZAPINE can cause serious side effects, including:

- **Changes in Feelings, Thoughts and Behaviour:** It is important that you have good communication with your healthcare professional about how you feel. Discussing your feelings and treatment with a friend or relative who can tell you if they think you are getting worse is also useful. Some patients may feel worse when they first start taking PRO-MIRTAZAPINE or when their dose is changed. You may have thoughts of hurting yourself or others. This can be more likely to happen if you have a history of suicidal thoughts or attempts and if you are aged 18 to 24 years old.
- **Agranulocytosis:** Severely low numbers of white blood cells that can lead to death.
- **Heart Problems:** PRO-MIRTAZAPINE can cause serious heart problems that can lead to death. There is a greater risk for these heart problems if you have or have a family history of heart rhythm problems (QT Prolongation).
- **Serotonin Syndrome and Neuroleptic Malignant Syndrome:** These are life-threatening conditions that affect your muscle tone and temperature control.
- **Serious Skin Problems:** PRO-MIRTAZAPINE can cause serious skin problems such as **Stevens-Johnson Syndrome (SJS), toxic epidermal necrolysis (TEN) and drug reaction with eosinophilia and systemic symptoms (DRESS)** that can be life-threatening.

See the “Serious side effects and what to do about them” table, below, for more information on these and other serious side effects.

Pregnancy: If you are pregnant, think you might be pregnant, or are thinking of becoming pregnant talk to your healthcare professional. If you get pregnant while taking PRO-MIRTAZAPINE you should discuss the risks and benefits of continuing your treatment with PRO-MIRTAZAPINE with your healthcare professional. Some newborns whose mothers took an SSRI or other newer antidepressants during pregnancy have developed complications at birth requiring prolonged hospitalization, breathing and feeding difficulties, seizures, tense or overly relaxed muscles, jitteriness and constant crying. In most cases, the newer antidepressant was taken during the third trimester of pregnancy. If your baby experiences any of these symptoms, contact your doctor as soon as you can.

Breastfeeding: If you are breastfeeding or thinking of breastfeeding discuss the risks and benefits of continuing your treatment with PRO-MIRTAZAPINE with your healthcare professional. PRO-MIRTAZAPINE passes into breastmilk. It is very important that you do NOT stop taking PRO-MIRTAZAPINE without first talking to your doctor.

Discontinuation Symptoms: Do not stop taking PRO-MIRTAZAPINE or change your dose without speaking to your healthcare professional. This can cause serious side effects. If you experience symptoms such as dizziness, abnormal dreams, numbness and tingling or electric shock sensations, agitation, anxiety, fatigue, confusion, headache, tremor, nausea,

vomiting or sweating when your dose has been lowered or changed, talk to your healthcare professional. These symptoms can also occur if you miss a dose.

Driving and using machines: Before you do tasks that require special attention, wait until you know how you respond to PRO-MIRTAZAPINE, it may affect your ability to be alert.

Blood tests: PRO-MIRTAZAPINE can cause abnormal blood test results. Your healthcare professional will decide when to perform blood tests and will interpret the results.

Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

Serious Drug Interactions

Do not use PRO-MIRTAZAPINE if you are taking or have recently taken:

- Monoamine Oxidase Inhibitors (such as; phenelzine, tranylcypromine, moclobemide, selegiline, linezolid, methylene blue)

The following may interact with PRO-MIRTAZAPINE:

- other antidepressants, such as Selective Serotonin Reuptake Inhibitors (SSRIs, i.e. paroxetine), venlafaxine, certain tricyclic antidepressants (i.e. amitriptyline), nefazodone, tryptophan
- ketoconazole; used to treat fungal infections
- cimetidine; used to treat reflux and stomach ulcers
- antibiotics used to treat bacterial infections, such as erythromycin and linezolid
- antiviral medicines used to treat Human Immunodeficiency Virus (HIV), such as a combination of fosamprenavir and ritonavir
- medicines used to treat epilepsy, such as carbamazepine and phenytoin
- rifampicin; an antibiotic used to treat tuberculosis
- warfarin; used to prevent blood clotting
- benzodiazepines; such as midazolam, oxazepam and diazepam
- medicines that may affect the heart's rhythm
- medicines used to treat mental health problems such as risperidone and lithium
- triptans; used to treat migraines
- opioids such as tramadol; used to treat moderate to severe pain
- St. John's Wort, an herbal product used to treat depression
- alcohol; you should not drink alcohol with you are taking PRO-MIRTAZAPINE

How to take PRO-MIRTAZAPINE:

- Take PRO-MIRTAZAPINE exactly as your healthcare professional tells you.
- Do not stop taking PRO-MIRTAZAPINE or change your dose without speaking to your healthcare professional.
- Take your tablet(s) at the same time each day, preferably in the evening before sleep.
- Swallow the tablets with water. Do not chew the tablets.

Usual adult dose:

15 mg – 45 mg once a day.

Overdose:

If you think you, or a person you are caring for, have taken too much PRO-MIRTAZAPINE, contact a healthcare professional, hospital emergency department or regional poison control centre immediately, even if there are no symptoms.

Signs that you have taken too much PRO-MIRTAZAPINE include; drowsiness, disorientation, increased heart rate. The symptoms of a possible overdose may include changes to your heart rhythm (fast, irregular heartbeat) and/or fainting which could be symptoms of a life-threatening condition known as Torsade de Pointes.

Missed Dose:

If you forget to take your evening dose, do not take the missed dose the next morning. Continue treatment in the evening (prior to sleep) with your normal dose. Do not take a double dose to make up for forgotten doses.

What are possible side effects from using PRO-MIRTAZAPINE?

These are not all the possible side effects you may feel when taking PRO-MIRTAZAPINE. If you experience any side effects not listed here, contact your healthcare professional.

Side effects may include:

- sleepiness, drowsiness
- weakness
- dry mouth
- increased appetite
- constipation
- weight gain
- dizziness
- headache
- muscle pain, back pain
- abnormal dreams, trouble sleeping, nightmares
- frequent urination
- itchiness, rash
- faintness (especially when you get up quickly from a lying or sitting position)
- urinary tract infections
- abnormal sensation in the skin (e.g., burning, stinging, tickling or tingly)

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
COMMON			
Amnesia: memory problems		✓	
RARE			
Agranulocytosis (severely low white blood cells): bruising, abnormal bleeding, signs of infection including fever, chills, sore throat, mouth sores/ulcers and swelling		✓	
Seizure (convulsion): muscle twitching, changes in emotions, confusion, loss of			✓

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
consciousness with uncontrollable shaking			
Hallucinations: hearing or seeing things that are not really there			✓
Mania: excessive happiness or irritability, racing thoughts, greatly increased energy, severe trouble sleeping, reckless behaviour			✓
Akathisia: feeling restless and unable to sit or stand still	✓		
Restless Legs: feeling of unrest during night in the legs, sudden muscle contractions in the legs	✓		
Liver problems: abdominal pain, yellowing of eyes or skin, dark urine, nausea, vomiting, loss of appetite			✓
Severe Skin Reactions; Stevens Johnson Syndrome (SJS), toxic epidermal necrolysis (TEN), Drug reaction with eosinophilia and systemic symptoms (DRESS): fever, severe rash, swollen lymph nodes, flu-like feeling, blisters and peeling of skin that may start in and around the mouth, nose, eyes and genitals and spread to other areas of the body, yellow skin or eyes, shortness of breath, dry cough, chest pain or discomfort, feeling thirsty, urinating less often, less urine, itching			✓
Hyponatremia (low sodium levels in the blood): feeling ill, weakness, drowsiness, confusion, achy, stiff or uncoordinated muscles			✓
Pancreatitis (inflammation of the pancreas): severe upper abdominal pain that lasts and gets worse when you lie down, nausea, vomiting, fever	✓		
VERY RARE			
Serotonin Syndrome and Neuroleptic Malignant Syndrome: agitation or restlessness, confusion, flushing, muscle twitching, tremor, involuntary eye movements, heavy sweating, high body temperature (>38°C), rigid muscles			✓
Changes in Feelings, Thoughts and Behaviour: anger, aggression, anxiety, thoughts of hurting yourself (suicide) or others			✓
UNKNOWN			
Heart Problems (QT Prolongation): abnormal heart rate or rhythm, palpitations,		✓	

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
dizziness, fainting			
Rhabdomyolysis: very dark “tea coloured” urine, muscle tenderness or weakness, muscle pain that you cannot explain		✓	
Hyperprolactinemia (high levels of prolactin in the blood): enlarged breasts and/or milky nipple discharge		✓	
Sleepwalking		✓	
Priapism: prolonged (longer than 4 hours) painful erection of the penis			✓

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, talk to your healthcare professional.

Reporting Side Effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

Storage:

- Store between 15°C and 30°C. Protect from light.
- Keep PRO-MIRTAZAPINE out of the reach and sight of children.
- Do not use PRO-MIRTAZAPINE after the expiry date indicated on the package.

If you want more information about PRO-MIRTAZAPINE:

- Talk to your healthcare professional
- Find the full Product Monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website (<https://www.canada.ca/en/health-canada/services/drugs-health-products/drug-products/drug-product-database.html>) or by contacting Pro Doc Ltée at 1-800-361-8559, www.prodoc.qc.ca or info@prodoc.qc.ca.

This leaflet was prepared by:

Pro Doc Ltée
Laval, Québec
H7L 3W9

Last revised: August 25, 2021